TIF® PROCEDURE FOR REFLUX

Is TIF Right For You?
Find Out Today
425-305-4688

The TIF procedure has helped thousands of people put a stop to reflux.

The TIF® procedure, the ***only*** solution that effectively treats reflux without incisions or metal implants.

Is TIF Right For You?

Fill Out the Form Below or Call To Request More Information or an Appointment

425-305-4688

First Name * | Last Name *

Phone # * | Zip Code *

Email *

REQUEST MORE INFORMATION OR AN APPOINTMENT

☑ I accept the terms of the Data Use Policy and acknowledge that I will receive communications via phone, email or text regarding my health or health risks, and/or more information on treatment options available in my area. View Data Use Policy

The TIF procedure delivers relief

- ✓ Minimally invasive: Fast recovery
- ✓ Restores valve to natural state: Maintain normal bodily functions
- ✓ No incisions: No scars
- ✓ No metal implant: No need to remove foreign object
- ✓ No dissection of stomach: No scar tissue

Incision-Free Surgery? Yes!

The TIF procedure takes advantage of advancements in minimally invasive procedures to restore your esophageal valve. TIF trained physicians use an endoscopic device that enables them to reach the valve and stomach without the need for incisions.

No More Reflux, No Incisions, No More PPIs.

20,000+ TIF procedures

FDA CLEARED — FDA cleared since 2007

81% of patients are off PPIs post-TIF procedure.[2] — 81% off daily PPIs[2]

Cleveland Providers

Dr. Leena Khaitan

216-532-1023

UH Digestive Health Institute
Dept of General Surgery
11100 Euclid Avenue
Cleveland, OH 44106
Driving Directions

UH Geauga Medical Center
Bariatrics Lab
13207 Ravenna Road
Chardon, OH 44024
Driving Directions

Dr. Jeffrey Marks

216-532-1027

UH Cleveland Medical Center
11100 Euclid Avenue
Cleveland, OH 44106
Driving Directions

UH Chagrin Highlands Health Center
909 Orange Place, Suite 3200
Orange Village, OH 44122
Driving Directions



Dr. John Rodriguez

216-532-1025

Lutheran Hospital - Cleveland Clinic
1730 W 25th Street, Suite 1E
Cleveland, OH 44113
Driving Directions

1. In the 2 studies in 56 patients where esophagitis was assessed at >=36 months, the median follow-up was 44 months, the weighted percentage of patients where the esophagus had completely healed was 83.94%.
2. In 11 TIF 2.0 on label studies collectively representing 568 patients where the follow-up was between 7 and 59 months, the weighted incidence percentage of patients completely off PPIs was 73.50%, and 7.91% of the patients reported occasional use of PPIs, for a total of 81.41% of the patients reporting no daily use of PPIs.

TIF® PROCEDURE FOR REFLUX

Is TIF Right For You?
Find Out Today
425-305-4688

Safety Disclaimer